

## **Nasco Agro Products**

**Address:** 53-55 Block-B, Road -3, Niketon, Gulshan-1, Dhaka

**Phone:** +880 1841-313036

**Email:** support@nascoagro.com

---

**Product Name:** HIPRAPOX ®

**Excerpt:** Live vaccine, fowl pox, in injectable freeze-dried tablet

**Category:** HIPRA

### **Description:**

#### **COMPOSITION:**

#### **PER DOSE:**

Attenuated live fowl pox virus, FPV-92 strain:  $10^4$ -  $10^{4,2}$  EID<sub>50</sub>

#### **INDICATIONS:**

**Chickens, Layers, Breeders and Turkeys:** Prevention of fowl pox.

#### **ADMINISTRATION ROUTE:**

Puncture in the wing membrane, or follicular scarification in the thigh.

- Reconstitute the freeze-dried tablet in the solvent included (1,000 doses).
- Shake gently until complete resuspension of the freeze-dried tablet before administration.
- Dip the two-pronged needle, which is included, in the vaccine solution.

**Chickens, Layers and Breeders:** Extend the wing membrane. Apply carefully the vaccine by puncturing the membrane in a manner that does not provoke lesions in the blood vessels. The needle must be freshly impregnated with the vaccine solution when performing each administration.

**Turkeys:** One drop of the vaccine solution will be applied by follicular scarification in the thigh.

#### **DOSAGE:**

**Chickens, Layers, Breeders and Turkeys:** 1 dose/bird.

- The veterinary surgeon will establish the most suitable vaccination programme according to the health conditions of each farm and area.
- After 7-10 days post-vaccination, one or two nodules should appear at the injection site which transform into scabs ("take"), indicating that the bird has been vaccinated. The scabs will disappear within 2-3 weeks after the vaccination.

#### **WITHDRAWAL TIME:**

0 days.

#### **SPECIAL PRECAUTIONS:**

- Store at +2 to +8 °C, protected from light.

#### **PACKAGING:**

Pack of 5 bottles of 1,000 doses + 5 bottles of solvent.

Under veterinary prescription